

January 2022

An Estimation of Blackbuck Abundance and Density in and around Jayamangali Blackbuck Conservation Reserve, Karnataka



© Hari Somashekhar

135, 14th Main, 30th Cross, Banashankari 2nd Stage,
Bengaluru - 560 070, India
TeleFax: +91-80-26716897
Website: www.holematthi.org

Office location:

Holématthi Nature Foundation
135, 14th Main, 30th Cross,
Banashankari 2nd Stage,
Bengaluru – 560 070
Karnataka, India
Telefax: +91 80 2671 6897
Email: malaika@ncf-india.org
Website: www.holematthi.org

Citation:

Gubbi, S., Mathew Chawla, M., Prabhu, K. & Suthar, S. (2022). An Estimation of Blackbuck Abundance and Density in and around Jayamangali Blackbuck Conservation Reserve, Karnataka, Holématthi Nature Foundation, Bengaluru, India.

Contents

Acknowledgements	01
Introduction	02
Methods	04
Results	09
Discussion	12
Conclusion and key management recommendations	14
References	15

Acknowledgements

We sincerely thank the Karnataka Forest Department for the permissions to carry out the study in Jayamangali Blackbuck Conservation Reserve, and for all the support provided by them. We are grateful to the British Asian Trust and the Karnataka Forest Department for funding this study.

We thank Dr S. Ramesh, Deputy Conservator of Forests, Tumkur Division, Tumkur; Mallikarjunappa H., Assistant Conservator of Forests, Madhugiri Sub-division, Madhugiri; Ravi C., Range Forest Officer, Madhugiri Range; and all Deputy Range Forest Officers, Forest Guards and Watchers who have proved to be the greatest source of assistance. We would like to acknowledge the support provided by DRFO Thippeswamy, Forest Guard Somappa V Ingu, and the Forest Watchers, Narasimhamurthi, Hanumantharayappa, Eeranna, Lakshmipathi and Nagaraju.

Introduction

The blackbuck (*Antilope cervicapra*) is a medium-sized antelope (31–45 kg) endemic to the Indian subcontinent. Blackbucks are primarily grazing herbivores that are representative species of semi-arid grasslands (Ranjitsinh, 1989). They also browse on leaf litter, flowers and fruits (Jhala, 1997; IUCN, 2017). Besides grasslands, they inhabit open scrub, open salt pans, open woodland and agricultural land (Ranjitsinh, 1989; Jhala, 1997). Blackbucks are group living animals; their social group size is variable and depends greatly on habitat openness (Isvaran, 2007). Larger group sizes in open or treeless habitats, such as grasslands, confer anti-predation benefits like early predator detection (Jarman, 1974; Isvaran, 2007). Blackbucks are key prey species for large carnivores in grasslands and other open habitats. Indian wolves (*Canis lupus pallipes*) are the main predators of blackbucks. Golden jackals (*Canis aureus*) and free-ranging dogs predate on blackbuck fawns (Jhala, 1993; Jethva & Jhala, 2004; Aiyadurai & Jhala, 2006; Jyoti & Rai, 2021). Blackbucks account for 12% of all wildlife species reportedly attacked by domestic dogs in India (Home *et al.*, 2017).

Historically, the blackbuck range encompassed most of the Indian subcontinent, south of the Himalayas (Ranjitsinh, 1989; IUCN, 2017). The species is now extinct in Pakistan and Bangladesh. In Nepal, the species went extinct in the 1970s, owing to excessive poaching. However, rediscovery of blackbuck populations in the region prompted several government-mediated interventions that led to the recovery of blackbuck populations in Nepal (Bist *et al.*, 2021). It is estimated that blackbuck population in Nepal is as low as 200 individuals (Bashishtha *et al.*, 2012). In India, the adult blackbuck population is estimated to be about 35,000 (IUCN, 2017). Given their locally abundant populations in India, blackbucks are currently listed as “Least Concern” in the IUCN Red List of Threatened Species. This is in stark comparison with the first IUCN assessment of the species, three decades ago, which suggested that populations were at a high risk of facing extinction in the wild (IUCN, 2017). The Government of India accords the blackbuck with the highest level of protection, under Schedule I of the Wildlife Protection Act of 1972.

In Karnataka, blackbucks occur mainly in the drier parts of the state (IUCN, 2017). Here, two protected areas have been notified chiefly for the conservation of the blackbuck: Ranibennur Blackbuck Sanctuary and Jayamangali Blackbuck Conservation Reserve. State-wide estimates of blackbuck populations in India do not exist. A few studies have estimated populations in protected areas, and even fewer studies, outside protected areas. A study by Sagar and Antoney (2017) provides a population estimate for Basur Amruth Mahal Kaval Conservation Reserve in Chikkamagaluru district using line transect sampling. They estimated 193 (c. 148–238) blackbucks to occur in an area of 7.36 km². This is the first study from Karnataka to provide blackbuck population estimates while accounting for imperfect detection.

From other parts of the country, robust population estimates are available from protected areas of Tamil Nadu, notably from Guindy National Park and Point Calimere Wildlife Sanctuary (Baskaran *et al.*, 2016; Arandhara *et al.*, 2020). Total count population surveys

which do not account for detection probability, have been carried out in Velavadar National Park, Gujarat (Jhala & Isvaran, 2016) and in Aligarh, Uttar Pradesh (Ahamad *et al.*, 2021).

Semi-arid grassland ecosystems are typical blackbuck habitats, enabling the species to reach high densities (Jhala & Isvaran, 2016). Grasses make up a major portion of the blackbuck diet (Jhala, 1997) and also provide camouflage to calves during periods when they are most vulnerable to predation (Priyadarshini *et al.*, 2021). Blackbucks avoid wooded habitats and prefer short grass with height less than 50 cm (Jhala, 1991). Consequently, in a habitat mosaic, blackbucks are restricted to open grassland habitats (Baskaran *et al.*, 2016; Arandhara *et al.*, 2021). In these habitats, the spread of the invasive shrub *Prosopis juliflora* has emerged as a significant threat to the ecology of grasslands. A recent study from Tamil Nadu demonstrated that the extent of *Prosopis* woodland and shrub cover has a negative impact on blackbuck density (Arandhara *et al.*, 2021).

In addition to the threat of non-native woody plants replacing native grassland, blackbucks are also at risk of coming in direct conflict with humans. Much of India's open grassland ecosystems have undergone conversion into agricultural land (Tian *et al.*, 2014), resulting in grassland-dependent wild herbivores opting for foraging sources available in croplands (Jhala, 1993). Blackbucks feeding on agricultural crops have been reported through several parts of its range in India (Chauhan & Singh, 1990; Jhala, 1993; Bayani *et al.*, 2016; Debata, 2017; Bajwa & Chauhan, 2019; Rai & Jyoti, 2019). In Abohar Wildlife Sanctuary in Punjab, the extent of crop loss due to nilgai and blackbucks was found to be as high as 40% (Bajwa & Chauhan, 2019). In retaliation to extensive crop loss, agriculturalists may employ preventive methods that can prove highly fatal to blackbucks. Chauhan & Singh (1990) have reported mass blackbuck mortality as a result of high voltage electric fencing used by farmers in Haryana.

In landscapes with high human densities, small patches of protected areas can serve as important refuges for blackbuck persistence (Krishna *et al.*, 2016), provided that levels of human activity are low in these areas (Jha & Isvaran, 2021, preprint). Given the context of lack of scientific information on blackbuck populations, shrinking grassland habitats, and rising anthropogenic pressures, we present a baseline population estimate for blackbucks in and around Jayamangali Blackbuck Conservation Reserve (JBCR). We use a systematic line transect sampling approach to provide baseline abundance and density estimates. A robust population baseline enables managers to draw inferences on temporal population trends and evaluate the effectiveness of management strategies, by comparing it against current population estimates (e.g. Rasphone *et al.*, 2021).

Methods

Study area

Jayamangali Blackbuck Conservation Reserve, situated in the Tumkur district of Karnataka (Map 1) was declared a conservation reserve in the year 2007. The conservation reserve consists of thorny shrubland, Eucalyptus (*Eucalyptus spp.*), Acacia (*Acacia auriculiformis*) and *Hardwickia binata* plantations, and remnant grassland (Gubbi *et al.* 2017) (Figure 1). The 3.24 km² reserve is surrounded by an agricultural-grassland mosaic (Figure 2). The commercial crops grown in the area include groundnut, horsegram, green pigeon peas, broad beans, chilli, sweet corn, castor, cotton, paddy, ragi, jowar, areca nut, eggplant, okra, tomato, watermelon, and muskmelon. In recent years, vine and banana cultivation has begun in the landscape. In the years 2019 to 2021, 47 farmers around JBCR have received a total of ₹3,06,686 ex-gratia payment by the government on account of crop loss caused by blackbucks. The affected crops were groundnut, green chilli, jowar, pigeon peas, ragi and corn (KFD, 2022).

Mammalian carnivores found in JBCR include leopard (*Panthera pardus fusca*), sloth bear (*Melursus ursinus*), Indian fox (*Vulpes bengalensis*), golden jackal (*Canis aureus*), jungle cat (*Felis chaus*), rusty-spotted cat (*Prionailurus rubiginosus*), small Indian civet (*Viverricula indica*), and grey mongoose (*Herpestes edwardsii*) (Gubbi *et al.* 2017). The Indian grey wolf (*Canis lupus pallipes*) has been documented in the study region previously (Gubbi, pers. obs.); however there seems to be little evidence of its presence in the area in recent times.

Herbivores besides the blackbuck are chital (*Axis axis*), wild pig (*Sus scrofa*), black-naped hare (*Lepus nigricollis*) and Indian porcupine (*Hystrix indica*) (Gubbi *et al.* 2017).

Data collection

A recce was carried out in the month of December 2021 to obtain information on terrain type and accessibility of habitats (Figure 3). A buffer of 5 km radius from the centre of the conservation reserve was drawn to include both, the protected area and surrounding agricultural fields encompassing a diversity of potential blackbuck habitats (Figure 4). To create line transects, the buffer area was divided into grids of 1 x 1 km, using a sampling tool in ArcGIS 10.3 (Esri, 2011). Line transects were systematically placed 2 km apart from each other, at an angle of 0°, facing north (Map 1). The lines were separated by a 2 km distance to ensure that all detections made from the line are independent of each other (Buckland & Turnock, 1992). A total of 16 line transects of 3 km length were created, however, two lines were excluded due to water logging. The resulting study area covered 140 km² in area. After this, the start and end points of each line transect were identified using a GPS and compass; and vegetation was cleared along the lines to enable the observer to walk with ease. It was ensured that the width of the lines were such that animals were neither attracted nor distracted from the lines.


Data was collected from 10th January to 31st January 2022 and transect lines were walked during early morning (6.00-8.00 h) and late afternoons (16.00-18.00 h) when blackbucks are most active. Each transect was walked 6 times, to adequately sample the study area and to increase precision. Sampling was carried out by one experienced observer and a Forest Department staff member who walked along the transect line and recorded detected individuals or a group of individuals. At every detection, distance to the animal or to the centre of the animal group was recorded using the Bushnell G Force DX rangefinder; and the angle to the animal and transect line angle was measured with the Sunto MC-2 compass. In order to obtain robust density and abundance estimates, we ensured that all assumptions of the line transect sampling approach were met (Buckland *et al.*, 2001). Given that our team comprised of trained field biologists, errors in measurements were minimized.



Figure 1. Jayamangali Blackbuck Conservation Reserve is dominated by closed habitats comprising of *Hardwickia binata* plantations.



Figure 2. The area surrounding the reserve is an agriculture-grassland mosaic.



Map 1. The line transect sampling design created for this study to estimate blackbuck abundance in and around Jayamangali Blackbuck Conservation Reserve; the study area measures 140 km².



Figure 3. In the month of December 2021, a recce was conducted to ascertain the accessibility of transect lines.



Figure 4. Line transect sampling was carried out in natural habitats as well as the agricultural matrix surrounding the Jayamangali Blackbuck Conservation Reserve.



Figure 5. Blackbucks show sexual dimorphism; adult males have a dark pelage with spiralled horns.

Data analysis

The distance sampling analysis was carried out in RStudio (RStudio Team, 2021) using the package ‘Distance’ (Miller *et al.*, 2019). The main inputs for the analysis are perpendicular distances, sampling effort and cluster or group size. 5% of the data at the largest distances were truncated to exclude outliers, as recommended by Buckland *et al.*, 2010. Six detection function models were fitted to the data; goodness of fit and Akaike Information Criterion (AIC) scores were computed to select the model with the best fit.




Figure 6. Histogram of blackbuck sighting frequency as a function of perpendicular distance of detections from the transect line. The graph reveals some heaping (rounding) of untruncated data beyond 400 m.

Results

The total sampling effort was 214.8 km. A total of 784 blackbuck individuals were recorded and the mean group size was found to be 4.92. The number of blackbuck detections made at each transect is listed in Table 1 and visualized in Map 2. We selected the half-normal key function model as the model with the best fit, according to AIC scores and the goodness of fit plots. The average detection probability (\hat{p}) deduced by the model is 0.53 (± 0.03 SE) (Figure 7).

The abundance of blackbucks in the study area (140 km²) is 2,144.9 (± 848.9 SE) individuals. The density of blackbucks is 15.3 (± 6.0 SE) individuals/km² (Table 2).

Table 1. Key observations of the line transect sampling carried out to estimate blackbuck density and abundance in and around Jayamangali Blackbuck Conservation Reserve.

Transect No.	Transect length (km)	Sampling effort (km)	No. of detections	No. of individuals	Encounter rate
JM-01	2.1	12.6	4	48	0.32
JM-02	3	18	0	0	0
JM-03	2.5	15	7	40	0.47
JM-04	2.6	15.6	17	46	1.09
JM-06	2.1	12.6	0	0	0
JM-07	3	18	3	7	0.17
JM-08	3	18	81	338	4.5
JM-09	3	18	30	220	1.66
JM-10	3	18	18	84	4.66
JM-12	2.7	16.2	1	1	0.006
JM-13	2.6	15.6	0	0	0
JM-14	2.4	14.4	0	0	0
JM-15	1.9	11.4	0	0	0
JM-16	1.9	11.4	0	0	0

Table 2. Estimated blackbuck density in and around Jayamangali Blackbuck Conservation Reserve (140 km²).

Total sampling effort (km)	Number of detections (n)	Mean cluster size	Total encounter rate	Density of individuals \hat{d} (per km ²)	Coefficient of Variance (\hat{d}) (%)
214.8	161	4.92	0.92	15.32	39.58%





Figure 7. The detection probability curve of the half-normal key function model with 5% truncation of perpendicular distances.



Map 2. Blackbuck detections on different transect lines during the study in and around Jayamangali Blackbuck Conservation Reserve.

Discussion

We used a systematic line transect sampling approach to provide abundance and density estimates for blackbucks in and around JBCR, Karnataka. Our study provides the first scientific estimate of blackbuck populations from this area. In addition, this is the second study that has estimated blackbuck populations in Karnataka, while accounting for imperfect detection.

The density of blackbucks is lower compared to other areas where similar studies have been carried out (Table 3). However, the abundance estimated for the entire study area is high. Our study area extends beyond the conservation reserve into human use areas like agricultural fields. It was important to study the population over a larger area since JBCR itself is very small (3.24 km²) and blackbucks range beyond the conservation reserve boundaries. Additionally, we found that blackbuck detectability was very low inside the conservation reserve. JBCR is dominated by woody plantations and blackbucks have been reported to avoid wooded habitats (Arandhara *et al.*, 2021).

Table 3. A comparison of blackbuck densities between the current study and previous studies in southern India where similar scientific methods were used.

Study site	Vegetation types	Study area (km ²)	Blackbuck density per km ² (SE)
Basur Amruth Mahal Kaaval Conservation Reserve, Karnataka (Sagar & Antoney, 2017)	Open grassland, invasive thorny shrubland	7.6	26.23 (±6)
Point Calimere Wildlife Sanctuary, Tamil Nadu (Baskaran <i>et al.</i> , 2016)	Tropical dry evergreen forest, open grassland, coastal scrub, invasive thorny shrubland	30.0	50.2 (±4.79)
Point Calimere Wildlife Sanctuary, Tamil Nadu (Arandhara <i>et al.</i> , 2020)	Tropical dry evergreen forest, open grassland, coastal scrub, invasive thorny shrubland	26.5	27.9 (±3.3)
In and around Jayamangali Blackbuck Reserve, Karnataka (current study)	Shrubland, plantations, open grassland, agricultural crop lands	140.0	15.32 (±6.06)

Blackbucks are highly specialized to semi-arid, short-grass ecosystems (Jhala & Isvaran, 2016) and loss of natural grasslands is one of the key threats to blackbuck conservation (IUCN, 2017). Previous studies recommend maintaining open grassland habitats, by preventing the establishment and invasion of exotic woody plants (Jhala, 1993; Arandhara *et al.*, 2021). Afforestation in grassland habitats, particularly with exotic trees like *Eucalyptus*, significantly impacts biodiversity and ecosystem functioning (Cravino & Brazeiro, 2021). It is essential that future management plans for JBCR are rooted in a scientifically sound understanding of blackbuck ecology. We recommend the removal of existing woody plantations from JBCR and that no future plantations are undertaken. Woody plantations have possibly eliminated the Indian wolf and have encouraged the immigration of leopards into the area. Past camera trapping studies have recorded the presence of three leopard individuals in JBCR (Gubbi *et al.*, 2017). The possible local extinction of wolves in Jayamangali may have impacted blackbuck populations, since wolves are the main natural predators of blackbucks (Jhala, 1993). In the absence of a key predator, blackbuck populations could significantly rise. This can in turn aggravate human-blackbuck conflict instances in this agricultural landscape. While natural grassland predators are necessary for regulating blackbuck populations, predators in the form of free-ranging dogs are a threat to blackbucks in JBCR. They have been observed to hunt both fawn and adult blackbucks in the region (Gubbi, pers. obs.).

Robust population estimates are important to scientifically understand the effectiveness of protected areas in species conservation. We believe that the baseline results of this study have important conservation implications for blackbuck conservation in JBCR and across Karnataka. We recommend that annual blackbuck population estimates be conducted using distance sampling in JBCR. This will provide a temporal understanding of trends in blackbuck populations, and accordingly inform conservation and management plans.

Conclusion

It is estimated that the total blackbuck population spread across India and Nepal is 35,000. This makes JBCR a significant stronghold for blackbucks in Karnataka and India. This area is perhaps a key area for the long-term survival of the species in the state. However, the peaceful co-existence of people and blackbucks in JBCR will depend on the restoration of native grasslands and the effective mitigation of potential conflict with farmers in the landscape.

Key management recommendations

1. Removal of woody plantations such as *Eucalyptus* spp., *Acacia auriculiformis*, *Hardwickia binata*, and *Tamarindus indica* from JBCR.
2. Annual population monitoring using distance sampling by line transect methodology.
3. Dispersal of timely and appropriate ex-gratia payments to farmers during human-blackbuck conflict instances.
4. Providing suitable incentives to farmers to grow blackbuck-friendly crops (horsegram, groundnut).
5. Making additions to the eco-sensitive zone plan to account for changing cropping patterns around JBCR and to suggest suitable crop alternatives.
6. Controlling the free-ranging dog population, given their threat to blackbucks as predators.

References

- Ahamad, M., Khan, J. A. & Kumar, S. (2021). Blackbuck *Antilope cervicapra* (Mammalia: Cetartiodactyla: Bovidae) estimates in human-dominated landscape in Aligarh, Uttar Pradesh, India. *Journal of Threatened Taxa*, 13(9): 19232–19238.
- Aiyadurai, A. & Jhala, Y. V. (2006). Foraging and habitat use by golden jackals (*Canis aureus*) in the Bhal region, Gujarat, India. *Journal of Bombay Natural History Society*, 103:5–12.
- Arandhara, S., Sathishkumar, S. & Baskaran, N. (2020). Modelling the effect of covariates on the detectability and density of native blackbucks and invasive feral-horse using Multiple Covariate Distance Sampling at Point Calimere Wildlife Sanctuary, Southern India. *Mammalian Biology*, 100, 173–186.
- Arandhara, S., Sathishkumar, S., Gupta, S. & Baskaran, N (2021). Influence of invasive *Prosopis juliflora* on the distribution and ecology of native blackbuck in protected areas of Tamil Nadu, India. *European Journal of Wildlife Research*, 67(3): 48.
- Bajwa, P. & Chauhan, N. S. (2019). Assessment of crop damage caused by Asian antelopes compared to local people perception in the community conserved Abohar Wildlife Sanctuary, Northwestern India. *Écoscience*, 26(4).
- Bashistha, M., Neupane, B. K. & Khanal, S. N. (2012). *Antilope cervicapra* Blackbuck in Nepal: Population Status, Conservation and Translocation Issues of Blackbuck in the Blackbuck Conservation Area, Bardiya, Nepal. LAP Lambert Academic Publishing, Saarbrücken.
- Baskaran, N., Ramkumaran, K. & Karthikeyan, G. (2016). Spatial and Dietary Overlap between Blackbuck (*Antilope cervicapra*) and Feral Horse (*Equus caballus*) at Point Calimere Wildlife Sanctuary, Southern India: Competition between Native Versus Introduced Species. *Mammalian Biology*, 81, 295–302.
- Bist, B. S., Ghimire, P., Poudyal, L. P., Pokharel, C. P., Sharma, P. & Pathak, K. (2021). From extinction to recovery: the case of blackbuck *Antilope cervicapra* from Nepal. *Mammal Research*, 66, 519–523.
- Buckland, S. T., Anderson, D. R., Burnham, K. P., Laake, J. L.; Borchers, D. L. & Thomas, L. (2001). Introduction to Distance Sampling, Oxford University Press.
- Buckland, S. T. & Turnock, B. J. (1992). A Robust Line Transect Method. *Biometrics*, 48(3): 901–909.
- Chauhan, N.P.S. & Singh, R. (1990). Crop damage by overabundant populations of nilgai and blackbuck in Haryana (India) and its management. Proceedings of the Fourteenth Vertebrate Pest Conference. Red Lion Inn, Sacramento, California 13: 218–220.

- Cravino, A. & Brazeiro, A. (2021). Grassland afforestation in South America: Local scale impacts of eucalyptus plantations on Uruguayan mammals. *Forest Ecology and Management*, 484: 118937.
- Debata, S. (2017). Population size, herd structure and sex ratio of the Blackbuck (*Antelope cervicapra* Linn.) (Mammalia: Cetartiodactyla: Bovidae) in a human-dominated area in Odisha, India. *Journal of Threatened Taxa*, 9(11): 10953–10955.
- ESRI (2011). ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute.
- Gubbi, S., Nagashettihalli, H., Bhat, R., Poornesha, H.C., Anoop, A., & Madhusudan, M.D. (2017). Ecology and conservation of leopards in protected and multiple use forests in Karnataka. Nature Conservation Foundation, Mysore, India.
- Home, C., Bhatnagar, Y. V. & Vanak, A. T. (2017). Canine Conundrum: domestic dogs as an invasive species and their impacts on wildlife in India. *Animal Conservation*, 21(4): 275–282.
- Isvaran, K. (2007). Intraspecific variation in group size in the blackbuck antelope: The roles of habitat structure and forage at different spatial scales. *Oecologia*, 154(2): 435–444.
- IUCN SSC Antelope Specialist Group. (2017). *Antelope cervicapra*. The IUCN Red List of Threatened Species: e.T1681A50181949.
- Jarman, P. J. (1974). The Social Organisation of Antelope in Relation To Their Ecology. *Behaviour*, 48(1–4): (1215–267).
- Jethva, B.D. & Jhala, Y. V. (2004). Foraging ecology, economics and conservation of Indian wolves in the *Bhal* region of Gujarat, Western India. *Biological Conservation*, 116(3): 351–357.
- Jha, R. R. & Isvaran, K. (2021). Antelope space-use and behavioural responses to varying anthropogenic influences may permit long-term persistence in a highly human-dominated landscape. Preprint available at bioRxiv: <https://doi.org/10.1101/2021.09.15.45990>.
- Jhala, Y. V. (1991). Habitat and population dynamics of wolves and blackbucks in Velavadar National Park, Gujarat, India. PhD dissertation. Virginia Polytechnic Institute and State University, Virginia.
- Jhala, Y. V. (1993). Predation on Blackbuck by Wolves in Velavadar National Park, Gujarat, India. *Conservation Biology*, 7(4): 874–881.
- Jhala, Y. V. (1997). Seasonal Effects on the Nutritional Ecology of Blackbuck *Antelope cervicapra*. *Journal of Applied Ecology*, 34(6): 1348–1358.
- Jhala, Y. V. & Isvaran, K. (2016) Behavioural ecology of a grassland antelope, the blackbuck *Antelope cervicapra*: linking habitat, ecology and behaviour. In: The ecology of large herbivores in South and Southeast Asia. Springer, Dordrecht.

- Jyoti & Rai, D. (2021). Measures of Sociality, Social Organization and Population Structure in Blackbuck, *Antilope cervicapra* (Linnaeus, 1758). *Proceedings of the Zoological Society*, 74(3):268–279.
- Karnataka Forest Department (2022). Publicity and ICT. <https://aranya.gov.in/>.
- Krishna, Y. C., Kumar, A. & Isvaran, K. (2016). Wild Ungulate Decision-Making and the Role of Tiny Refuges in Human-Dominated Landscapes. *PLoS ONE*, 11(3): e0151748.
- Miller, D. L., Rexstad, E., Thomas, L., Marshall, L., Laake, J.L. (2019). Distance Sampling in R. *Journal of Statistical Software*, 89(1): 1–28.
- Rai, D. & Jyoti (2019). Crowding, group size and population structure of the Blackbuck *Antilope cervicapra* (Linnaeus, 1758) (Mammalia: Cetartiodactyla: Bovidae) in the semi-arid habitat of Haryana, India. *Journal of Threatened Taxa*, 11(9): 14194–14203. <https://doi.org/10.11609/jott.4788.11.9.14194-14203>.
- Ranjitsinh, M. K. (1989). The Indian Blackbuck. Natraj Publishers. Dehradun.
- Rasphone, A., Kamler, J. F., Tobler, M. & Macdonald, D. W. (2021). Density trends of wild felids in northern Laos. *Biodiversity Conservation*, 30, 1881–1897.
- RStudio Team (2021). RStudio: Integrated Development Environment for R. RStudio, PBC, Boston, MA. <http://www.rstudio.com/>.
- Priyadarshini, K. V. R., Gort, C., Rice, C. G., Yoganand, K. (2021). The reproductive phenology of blackbuck: influence of seasonal nutritional resources and flexible lactation as an adaptive strategy. *Journal of Zoology*, 316(1): 11–22.
- Sagar, H.S.C. & Antoney, P. U. (2017). Measuring Indian Blackbuck *Antilope cervicapra* (Mammalia: Cetartiodactyla: Bovidae) abundance at Basur Amruth Mahal Kaval Conservation Reserve, Chikkamagaluru, southern India. *Journal of Threatened Taxa*, 9(7): 10468–10472.
- Tian, H., Banger, K., Bo, T., & Dadhwal, V. K. (2014). History of land use in India during 1880–2010: Large-scale land transformations reconstructed from satellite data and historical archives. *Global and Planetary Change*, 121: 78–88.

Research Team

Dr. Sanjay Gubbi

Kiran Prabhu

Shravan Suthar

Malaika Mathew Chawla

Poornesha H. C.

Sandesh Appu Naik

Ruma K. Kandurkar

Praveen T. V.